



# SOFT MATTER INTERFACES (SMI)

**Group Leader:** Elaine DiMasi<sup>1</sup> **Proposal Team:** E. DiMasi<sup>1</sup>, A. Fluerasu<sup>1</sup>, M. Fukuto<sup>1</sup>, O. Gang<sup>1</sup>, B. Hsiao<sup>2</sup>, J. Kline<sup>3</sup>, T. Koga<sup>2</sup>, I. Kuzmenko<sup>4</sup>, **B. Ocko**<sup>1</sup>, R. Pindak<sup>1</sup>, M. Schlossman<sup>4</sup>, H. Strey<sup>2</sup>, D. Vaknin<sup>5</sup>, L. Wiegart<sup>1</sup>, T. Xu<sup>6</sup>, K. Yager<sup>1</sup>, L. Yang<sup>1</sup>, J. Zhang<sup>4</sup>  
<sup>1</sup>Brookhaven National Laboratory, <sup>2</sup>Stony Brook University, <sup>3</sup>NIST, <sup>4</sup>Argonne National Laboratory, <sup>5</sup>University of Illinois, <sup>6</sup>Ames Laboratory, <sup>7</sup>University of California, Berkeley




## TECHNIQUES AND CAPABILITIES

- Enables in-situ studies of soft-matter interface structures using specialized surface sensitive x-ray scattering techniques.
- Two dedicated endstations will allow studies to be carried out at the solid/vapor, solid/liquid, liquid/vapor, liquid/solid and liquid/liquid interfaces, with new tender energy capabilities.
- The high brightness of NSLS-II will enable microbeam modes, high  $q$  resolution, time resolved measurements, and a wide  $q$  range for simultaneous data acquisition.
- Simultaneous GISAXS / GIWAXS / GIXD
- Diffuse scattering, bragg rods
- Fast x-ray reflectivity from liquid interfaces
- Predicting  $10^{15}$  ph/s from undulator source
- Energy range 2 to 24 keV
- Energy tunability for resonant studies
- Beam size  $5\text{-}500\ \mu\text{m} \times 2\text{-}100\ \mu\text{m}$  ( $h \times v$ )
- Configurable horizontal & vertical focusing



## APPLICATIONS in ENERGY, ENVIRONMENT, and BIOMOLECULAR MATERIALS



### ENERGY

Microphase behavior and molecular orientations in thin film organic photovoltaic devices and organic LEDs, promising new classes of energy materials; gas hydrate formation at the gas-water interface; structure-function relation in device materials

### ENVIRONMENT


Extractant-mediated transfer of metal ions from aqueous to immiscible organic phase; carbonate mineralization for carbon sequestration

### BIOMOLECULAR MATERIALS

Fundamental understanding of self-assembly to create new functional bio, nano, hybrid, and hierarchical materials

## INSTRUMENTATION ADVANCES

- Windowless path 2-4 keV for resonance from P, S, Cl, K, and Ca for biomolecules
- Simultaneous small and wide angle scattering with  $> 200\text{Hz}$  time resolution and new data visualization
- New capabilities for Liquids with tender x-rays and fast reflectivity



Poster created by Lutz Wiegart, Interim Project Leader, 2011.  
 Updated by SMI Beamline Development Group,  
 Elaine DiMasi, Warren Halbig, and Amanda King, 2012.

